

FOMIN–KIRILLOV ALGEBRAS

LEANDRO VENDRAMIN

ABSTRACT. This is an extended abstract of the talk given in the Oberwolfach miniworkshop “Nichols algebras and Weyl groupoids” in October 2012.

For an integer $n \geq 3$ denote by $\mathcal{E}_n$ the algebra (of type A_{n-1}) with generators $x_{(ij)}$, where $1 \leq i < j \leq n$, and relations

$$\begin{aligned} x_{(ij)}^2 &= 0, & \text{for } 1 \leq i < j \leq n, \\ x_{(ij)}x_{(jk)} &= x_{(jk)}x_{(ik)} + x_{(ik)}x_{(ij)}, & \text{for } 1 \leq i < j < k \leq n, \\ x_{(jk)}x_{(ij)} &= x_{(ik)}x_{(jk)} + x_{(ij)}x_{(ik)}, & \text{for } 1 \leq i < j < k \leq n, \\ x_{(ij)}x_{(kl)} &= x_{(kl)}x_{(ij)}, & \text{for any discinct } i, j, k, l. \end{aligned}$$

The algebras $\mathcal{E}_n$ are graded by $\deg(x_{(ij)}) = 1$. Of course, it is natural to ask if $\mathcal{E}_n$ is finite-dimensional. It is known that $\mathcal{E}_n$ is finite-dimensional if $n \leq 5$. It has been conjectured that $\dim \mathcal{E}_n = \infty$ for $n \geq 6$.

Example 1. The algebra $\mathcal{E}_3$ has dimension 12. The Hilbert series $\mathcal{H}_3(t)$ of $\mathcal{E}_3$ is a polynomial of degree 4: $\mathcal{H}_3(t) = (2)_t^2(3)_t$, where $(k)_t = 1 + t + \dots + t^{k-1}$.

Example 2. Computer calculations yield $\dim \mathcal{E}_4 = 576$. The Hilbert series $\mathcal{H}_4(t)$ of $\mathcal{E}_4$ is a polynomial of degree 12: $\mathcal{H}_4(t) = (2)_t^2(3)_t^2(4)_t^2$.

Example 3. Computer calculations yield $\dim \mathcal{E}_5 = 8294400$. The Hilbert series $\mathcal{H}_5(t)$ of $\mathcal{E}_5$ is a polynomial of degree 40: $\mathcal{H}_5(t) = (4)_t^4(5)_t^2(6)_t^4$.

Example 4. The Hilbert series $\mathcal{H}_6(t)$ of $\mathcal{E}_6$ cannot be written as a product of t -numbers. Further,

$$\mathcal{H}_6(t) = 1 + 15t + 125t^2 + 765t^3 + 3831t^4 + 16605t^5 + 64432t^6 + 228855t^7 + \dots$$

In [3], Fomin and Kirillov introduced the algebras $\mathcal{E}_n$ as a new model for the Schubert calculus of a flag manifold. They proved that $\mathcal{E}_n$ contains a commutative subalgebra isomorphic the cohomology ring of the flag manifold. In [1], Bazlov proved that Nichols algebras provide the correct setting for this model of Schubert calculus. But what is the relation between the algebras $\mathcal{E}_n$ and Nichols algebras?

Let V_n be the vector space with basis $\{v_{(ij)} \mid 1 \leq i < j \leq n\}$ and consider the map $c \in \mathbf{GL}(V_n \otimes V_n)$ defined by

$$c(v_\sigma \otimes v_\tau) = \chi(\sigma, \tau) v_{\sigma\tau\sigma^{-1}} \otimes v_\sigma, \quad \chi(\sigma, \tau) = \begin{cases} 1 & \text{if } \sigma(i) < \sigma(j), \\ -1 & \text{otherwise,} \end{cases}$$

where σ and τ are transpositions, and $\tau = (i \ j)$ with $i < j$. Since (V_n, c) is a braided vector space, it is possible to consider the Nichols algebra $\mathfrak{B}(V_n)$. Bazlov proved

This work was supported by Conicet and the Alexander von Humboldt Foundation. The author thanks Bernard Leclerc for conversations about cluster algebras and Lie theory.

that $\mathfrak{B}(V_n)$ contains a commutative subalgebra isomorphic to the cohomology ring of the flag manifold.

It is known that $\mathfrak{B}(V_n) = \mathcal{E}_n$ if $3 \leq n \leq 5$; this was proved by Milinski and Schneider for $n \leq 4$, and by Graña for $n = 5$. It has been conjectured that $\mathfrak{B}(V_n)$ is quadratic and $\mathfrak{B}(V_n) = \mathcal{E}_n$; see for example [5] and [1].

There are many others interesting conjectures about Fomin–Kirillov algebras. In [5], Majid wrote that it might be possible to find a relation between Fomin–Kirillov algebras and the representation theory of preprojective algebras.

Let Λ be the preprojective algebra of a quiver of type A_{n-1} . It is known that the number of indecomposable modules over Λ is 4 if $n = 3$, 12 if $n = 4$, and 40 if $n = 5$. Further, Λ is of infinite representation type if $n \geq 6$. Majid noticed that the number of indecomposable modules over Λ is equal to the degree of the Hilbert series of $\mathcal{E}_n$, at least for $3 \leq n \leq 5$. Majid’s conjecture does not have a precise formulation, but it states that this numerology is not an accident.

To conclude, we restate Majid’s observation in terms of cluster algebras. Let $n \geq 2$, $G = \mathbf{SL}_n$, and N be the subgroup of upper triangular matrices with ones in the diagonal. In [2], Berenstein, Fomin and Zelevinski proved that $\mathbb{C}[N]$, the coordinate ring of N , has a cluster algebra structure. Furthermore, the number of clusters of $\mathbb{C}[N]$ is given by the following table:

Lie type of G	Number of clusters
A_2	4
A_3	12
A_4	40
others	∞

Geiss, Leclerc and Schröer established a relation between the number of clusters of $\mathbb{C}[N]$ and the number of indecomposable modules over the preprojective algebra Λ , see for example [4]. This implies that Majid’s observation can be translated into the combinatorial language of cluster algebras.

REFERENCES

- [1] Y. Bazlov. Nichols-Woronowicz algebra model for Schubert calculus on Coxeter groups. *J. Algebra*, 297(2):372–399, 2006.
- [2] A. Berenstein, S. Fomin, and A. Zelevinsky. Cluster algebras. III. Upper bounds and double Bruhat cells. *Duke Math. J.*, 126(1):1–52, 2005.
- [3] S. Fomin and A. N. Kirillov. Quadratic algebras, Dunkl elements, and Schubert calculus. In *Advances in geometry*, volume 172 of *Progr. Math.*, pages 147–182. Birkhäuser Boston, Boston, MA, 1999.
- [4] B. Leclerc. Cluster algebras and representation theory. In *Proceedings of the International Congress of Mathematicians. Volume IV*, pages 2471–2488, New Delhi, 2010. Hindustan Book Agency.
- [5] S. Majid. Noncommutative differentials and Yang-Mills on permutation groups S_n . In *Hopf algebras in noncommutative geometry and physics*, volume 239 of *Lecture Notes in Pure and Appl. Math.*, pages 189–213. Dekker, New York, 2005.

PHILIPPS-UNIVERSITÄT MARBURG, FB MATHEMATIK UND INFORMATIK, HANS-MEERWEIN-STRASSE, 35032 MARBURG, GERMANY
E-mail address: lvendramin@dm.uba.ar